





Antimicrobial Use and Resistance (AUR) Option: Protocols and Definitions

Mary Andrus, BA, RN, CIC

Division of Healthcare Quality Promotion

SAFER • HEALTHIER • PEOPLE™

Target Audience

- This training session is designed for those who will collect and analyze Pharmacy or Microbiology data in the Antimicrobial Use and Resistance (AUR) option of the Medication-associated module of the Patient Safety component of NHSN. This may include:
 - NHSN Facility Administrator
 - Patient Safety Primary Contact
 - Infection Control Professional (ICP)
 - Epidemiologist
 - Microbiologist
 - Pharmacist
 - Data entry staff

Objectives

- Outline the impact of antimicrobial use and resistance in healthcare
- Identify the structure and purpose of the AUR option within the Patient Safety Component of NHSN
- 3. Review the requirements for participation in the AUR option
- 4. Define location types required in the AUR option


Objectives (cont.)

- 5. Describe AUR elements in
 - Monthly Reporting Plan
 - Summary data for Microbiology
 - Summary data for Pharmacy
- Demonstrate the use of AUR rates to improve infection control practices and antimicrobial prescription practices

Background

- Antimicrobial-resistant organisms are increasing rapidly in the U.S.
- Reasons:
 - Transmission between patients
 - Selection of resistant organisms because of antimicrobial receipt






http://www.cdc.gov/ncidod/dhqp/pdf/ar/mdroGuideline2006.pdf

AUR Option Purpose

- Assist hospitals in collecting data on antimicrobial use or bacterial resistance to antibiotics or both
- Feedback rates of antimicrobial use and resistance to encourage appropriate antimicrobial prescribing
- Encourage activities targeted toward the prevention of patient-to-patient transmission



Medicationassociated Module

Antimicrobial Use and Resistance Option

AUR - Microbiology

AUR - Pharmacy

Requirements

- Collect data for a minimum of 6 months* per calendar year
- An acceptable month of data includes:
 - Clinical isolates from microbiology or...
 - Antimicrobial agents used from pharmacy or...
 - Both
- Data must be submitted for specific location types within your hospital

^{*}Submission of fewer than 6 months will not be adequate to accurately measure antimicrobial resistance or use rates

Important Note!



- AUR Option does not collect data about patient infections
- Simultaneous collection of data using the Device-associated Event Module is strongly encouraged for the same months and in the same locations as followed in the AUR Option.

Monthly Reporting Plan - AUR

- A Plan must be submitted for AUR for each month
- The Plan must include all of the following locations:
 - At least one ICU or SCA
 - All non-ICU/SCA areas combined
 - All outpatient areas combined*
- For each above area, the facility chooses to monitor either microbiology data or pharmacy data* or both

* **Exception** — No pharmacy data are collected in outpatient areas!

AUR Location Categories

Intensive Care Unit (ICU): a patient care area that provides intensive observation, diagnosis, and therapeutic procedures for critically ill patients. Excludes step-down, intermediate, or telemetry care.

AUR Location Categories

- Specialty Care Area (SCA): a patient care area in which 80% of patients are of the following types:
 - Bone marrow/stem cell transplant
 - Solid organ transplant
 - Patients with hematologic or oncologic malignancies
 - Peritoneal or hemodialysis
 - Long term acute care

Location Definitions for AUR


- All Inpatient (non-ICU/SCA): a designated patient care location category for combining data from all inpatient areas that are not ICU or SCA.
 - The patients in this location are NHSN inpatients (i.e., those patients whose date of admission and discharge are different calendar dates)
 - Data reported as a single entity in AUR option


Location Definitions for AUR


- All Outpatient: a designated patient care location category which combines the data from all outpatient areas together
 - Patients are admitted and discharged on the same calendar date
 - Data from all outpatient areas combined and reported as a single entity in AUR option
 - Pharmacy data are <u>not</u> reported for outpatients in AUR





- AUR Locations are established during the Set Up process
- December 7 training "Facility Start Up"






















AUR – Microbiology Data



Microbiology Data Collection form

Antimicrobial Use and Resistance (AUR) Microbiology Laboratory Data - Monthly Report Form


OMB No. 0920-0666 Exp. Date: 02-29-2000

required for saving

*Facility ID#:	*Month: Oct	*Year: 2006	*Location 0	ode: ALLIN

	Susceptible*	Intermediate*	Resistant*	Total Tested*
GRAM POSITIVE ORGANIS	MS			
Coagulase-negative staphylo	cocci			
vancomycin				
Enterococcus spp.				
vancomycin				
Staphylococcus aureus		4	· · · · · · · · · · · · · · · · · · ·	
oxacillin				
vancomycin				1
GRAM NEGATIVE ORGANIS	MS			
Acinetobacter spp.				
amikacin				
cefepime				
ceftazidime				
imipenem				
piperacillin/tazobactam				
ampicillin/sulbactam	s			
Enterobacter spp.				
cefotaxime				
ceftazidime				
imipenem				
meropenem				
Escherichia coli	¥3			
cefotaxime				
ceftazidime				
ciprofloxacin				
imipenem				





- Report antimicrobial susceptibility test*
 results on all nonduplicate, clinical isolates
 processed by the laboratory during each
 month of monitoring
 - S Susceptible
 - I Intermediate
 - R Resistant

* Defined by Clinical Laboratory Standards Institute (CLSI)


Microbiology – Duplicate Isolates


- Do not include duplicate isolates.
 - A duplicate isolate is defined as an isolate of the same bacteria with the same antimicrobial susceptibility pattern in the same patient, regardless of specimen site, during a given calendar month





Microbiology – Surveillance Cultures

- Do not include surveillance cultures.
 - Surveillance cultures are those cultures reported as part of infection control surveillance such as stool cultures for vancomycin resistant enterococci (VRE) or routine nares screening for MRSA on admission













AUR – Microbiology Data

Number of resistant isolates

X 100

Resistance Rate =

Number of isolates tested



SAFER · HEALTHIER · PEOPLE™





AUR – Pharmacy Data



SAFER · HEALTHIER · PEOPLE™

Pharmacy Data Collection Form



cefadroxil

Antimicrobial Use and Resistance (AUR)


OMB No. 0920-0666 Exp. Date: 02-29-2008

Pharmacy Data - Monthly Report Form * required for *Year: 2006 *Location Code: RMT *Month: Oct *Patient Days: 633 *Facility ID#: **Parenteral Antibiotics** Antibiotic *Ouantity Used[†] Antibiotic *Ouantity Used[†] amikacin gentamicin ampicillin imipenem ampicillin**/sulbactam levofloxacin azithromycin linezolid aztreonam meropenem cefamandole metronidazole cefazolin moxifloxacin cefepime nafcillin cefmetazole ofloxacin cefotaxime oxacillin cefotetan penicillin G mill. I.U. cefoxitin pen. G benzathine mill. I.U ceftazidime procaine pen. G mill, I.U. ceftizoxime piperacillin ceftriaxone piperacillin**/tazobactam cefuroxime quinupristin **/dalfopristin cephalothin ticarcillin ciprofloxacin ticarcillin**/clavulanic acid clindamycin tigecycline daptomycin tobramycin trimethoprim**/sulfamethoxazole ertapenem erythromycin vancomycin gatifloxacin **Oral Antibiotics** Antibiotic *Quantity Used[†] Antibiotic *Quantity Used[†] amoxicillin gatifloxacin amoxicillin**/clavulanic acid levofloxacin linezolid ampicillin azithromycin lomefloxacin cefaclor metronidazole

a moxifloxacin




- The number of grams or million international units (mill.I.U.) are reported monthly for inpatients
- Selected oral and parenteral antimicrobial agents





Department of Health and Human Services Centers for Disease Control and Prevention





Department of Health and Human Services Centers for Disease Control and Prevention

	NHSN - National Healthcare Safety Network		NHSN Home	My Info Contact us Help
3N Home ting Plan it dure lary Data	Logged into Medical Center East (ID 10000 Facility Medical Center East (ID 10000) is fo Antim	ollowing PS component. icrobial Use a	ind Resistance (A acy Data	UR)
i sis y y ut	Mandatory fields marked with * Facility ID*: 10000 (Medic Location Code*: SICU-SURG Month*: October Year*: 2006 Total Patient Days*: 741	Enter the	number of patient days for or the month	Print PDF
	The state of the s			
	Parenteral Antibiotics		25/12	
	Parenteral Antibiotics Antibiotic	Quantity Used*	Antibiotic	Quantity Use
		Quantity Used*	Antibiotic GATI - Gatifloxacin	Quantity Use
	Antibiotic			
	Antibiotic AMK - Amikacin	g g	GATI - Gatifloxacin	g
	Antibiotic AMK - Amikacin AMP - Ampicillin	g g	GATI - Gatifloxacin GENT - Gentamicin	g g
	Antibiotic AMK - Amikacin AMP - Ampicillin AMPSUL - Ampicillin/sulbactam	g g g g g	GATI - Gatifloxacin GENT - Gentamicin IMI - Imipenem	g g
e	Antibiotic AMK - Amikacin AMP - Ampicillin AMPSUL - Ampicillin/sulbactam AZITH - Azithromycin	g g g g g	GATI - Gatifloxacin GENT - Gentamicin IMI - Imipenem LEVO - Levofloxacin	g g g g g



Department of Health and Human Services Centers for Disease Control and <u>Prevention</u>

NHSN - National Healthcare Safety Network | NHSN Home | My Info | Contact us | Help | Logged into Medical Center East (ID 10000) as MVA. 3N Home Facility Medical Center East (ID 10000) is following PS component. ting Plan Antimicrobial Use and Resistance (AUR) **Pharmacy Data** dure ary Data Mandatory fields marked with * Print PDF Facility ID*: 10000 (Medical Center East) sis Location Code*: SICU-SURGICALICU Month*: October Year*: 2006 ▼ Total Patient Days*: 741 **Parenteral Antibiotics** Quantity Used* **Antibiotic** Quantity Use **Antibiotic** GATI - Gatifloxacin AMK - Amikacin 31.0 g 0.0 AMP - Ampicillin 1338.0 GENT - Gentamicin 35.88 AMPSUL - Ampicillin/sulbactam **1866.0** 9 IMI - Imipenem 169.5 **78.5** g LEVO - Levofloxacin AZITH - Azithromycin 478.5

e



AZT - Aztreonam



44.0

LNZ - Linezolid




60.0 9
Local intranet

CEFEP - Cefepime	0.0	g	NAF - Nafcillin	825.0	g
CEFMET - Cefmetazole	0.0	g	OFLOX - Ofloxacin	0.0	g
CEFOT - Cefotaxime	0.0	g	OX - Oxacillin	0.0	g
CTET - Cefotetan	2.0	g	PENG - Penicillin G	5922.0	mill. I.
CEFOX - Cefoxitin	0.0	g	PENGBZ - Pen. G benzathine	50.8	mill. I.
CEFTAZ – Ceftazidime	477.0	g	PROPG - Procaine pen. G	0.0	mill. I.
CEFTIZ - Ceftizoxime	2.0	g	PIP - Piperacillin	1082.0	g
CEFTRX - Ceftriaxone	riaxone PIPTAZ - Piperacillin/tazobactam		PIPTAZ - Piperacillin/tazobactam	10148.0	g
CEFUR - Cefuroxime	79.5	g	QUIDAL - Quinupristin/dalfopristin	0.0	g
CEPH - Cephalothin	0.0	g	TICAR - Ticarcillin	0.0	g
CIPRO – Ciprofloxacin	8.4	g	TICLAV - Ticarcillin/clavulanic acid	90.0	g
CLIND - Clindamycin	3026.55	g	TOBRA - Tobramycin	12.56	g
DAPTO - Daptomycin	0.0	g	TMZ - Trimethoprim/sulfamethoxazole	99.2	g
ERTA - Ertapenem	18.0	g	VANC - Vancomycin	4507.3	g
ERYTH - Erythromycin	28.0	g			
Oral Antibiotics	140			test	
Antibiotic	Quantity	/ Used*	Antibiotic	Quantity	Used*
AMOX - Amoxacillin	50.5 g		GATI - Gatifloxacin	0.0	g
AMXCLV - Amoxacillin/clavulanic acid	moxacillin/clavulanic acid 162.275 g LEVO - Levofloxacin		LEVO - Levofloxacin	600.5	g
AMP - Ampicillin	47.0	g	LNZ - Linezolid	77.5	g

Local intranet

ERYTH - Erythromycın		g			
Oral Antibiotics					
Antibiotic	Quantity	y Used*	Antibiotic	Quantity	y U
AMOX - Amoxacillin	50.5	g	GATI - Gatifloxacin	47.0	g
AMXCLV - Amoxacillin/clavulanic acid	162.275	g	LEVO - Levofloxacin	600.5	g
AMP - Ampicillin	47.0	g	LNZ - Linezolid	77.5	g
AZITH - Azithromycin	112.8	g	LOMEF - Lomefloxacin	0.0	g
CEFAC - Cefaclor	151.0	g	METRO - Metronidazole	717.5	g
CEFAD - Cefadroxil	0.0	g	MOXI - Moxifloxacin	0.0	g
CEFIX - Cefixime	0.0	g	NOR - Norfloxacin	0.0	g
CEFPRO - Cefprozil	0.0	g	OFLOX - Ofloxacin	0.0	g
CEPHLX - Cephalexin	200.50	g	PENV - Penicillin V	5.5	g
CIPRO - Ciprofloxacin	83.5	g	SPAR - Sparfloxacin	0.0	g
CLARTH - Clarithromycin	181.25	g	TELITH - Telithromycin	0.0	g
CLIND - Clindamycin	65.60	g	TETRA - Tetracycline	199.0	g
DICLOX - Dicloxacillin	25.5	g	TMZ - Trimethoprim/sulfamethoxazole	381.526	g
DOXY - Doxycycline	190.3	g	VANC - Vancomycin	0.0	g
ERYTH - Erythromycin	45.75	g			





Calculating Antimicrobial Use Rates

DDD of antimicrobial =

Grams of antimicrobial used

DDD in grams*

* See Table 15 in NHSN Users Manual http://www.cdc.gov/ncidod/hip/nhsn/member s/PSProtocolsMay06.pdf



Class	Group	Antimicrobial Agent	DDD
3-lactams	Penicillin group	Penicillin G	1.2 x 10 ⁶ U*
		Procaine Penicillin G	$2.4 \times 10^6 U^*$
		Penicillin G benzathine	$1.2 \times 10^6 \mathrm{U}^*$
		Penicillin V	1 g*
	Ampicillin group	Ampicillin (parenteral)	2g
		Ampicillin (oral)	2g
		Ampicillin/sulbactam	2g
		Amoxicillin (oral)	1g
		Amoxicillin/Clavulanic Acid (oral)	1g
	Antistaphylococcal penicillins (Methicillin	Nafcillin	4g*
	group)	Oxacillin	2g
		Dicloxacillin (oral)	2g
	Antipseudomonal	Piperacillin	14g
	penicillins	Piperacillin/Tazobactam	14g
	•	Ticarcillin	15g
		Ticarcillin/Clavulanic Acid	15g
	1st-Generation	Cefazolin	3g
	cephalosporins	Cephalothin	4g
		Cefadroxil (oral)	2g
		Cephalexin (oral)	2g
	2nd-Generation	Cefotetan	4g
	cephalosporins	Cefmetazole	4g*
		Cefoxitin	6g
		Cefuroxime	3g
		Cefuroxime axetil (oral)	1g*
		Cefaclor (oral)	1g
		Cefprozil (oral)	1g
	3rd-Generation	Cefotaxime	4g
	cephalosporins	Ceftazidime	4g
		Ceftizoxime	4g
		Ceftriaxone	2g
		Cefixime (oral)	0.4g
		Cefipime	2g
	Carbapenems	Meropenem	2g
	-	Imipenem cilastatin	2g







Using AUR Rates to Improve Practices




Table 10. Pooled means and percentiles of the distribution of antimicrobial resistance rates*, by all ICUs combined, non-ICU inpatient units and by outpatients, ICARE/AUR, January 1998 through June 2004

All ICUs comb	oined					Percentile		
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%
MRSA	157	22,899	52.90	20.0	32.7	48.1	60.3	67.9
Methicillin-resistant CNS	141	13,553	76.60	57.0	69.4	76.3	83.8	88.4
Vancomycin-resistant Enterococcus spp	140	14,140	13.90	0	5	13.6	24.3	39.2
Ciprofloxacin/ofloxacin-resistant Pseudomonas aeruginosa	134	13,473	34.80	8.3	17.4	29.3	41.3	51.6
Levofloxacin-resistant P aeruginosa	68	5895	35.30	9.7	18.2	29.1	40.8	47.7
Imipenem-resistant P aeruginosa	123	11,986	19.10	4.8	8.3	13.2	25.5	38
Ceftaz idi me-re isistan t P aeruginosa	129	12,805	13.90	0	5	10.8	16.9	23.6
Piperacillin-resistant P aeruginosa	118	11,640	17.50	2.4	7.5	14.3	19.5	31.4
Cef3-resistant Enterobacter spp	111	5328	27.70	10.0	17.4	26.1	36.4	47.4
Carbapenem-resistant Enterobacter spp	93	4663	0.70	0	0	0	0	3.8
Cef3-resistant Klebsiella pneumoniae	119	7529	6.20	0	0	2.0	8.0	20.7
Cef3-resistant Escherichia coli	140	12,011	1.30	0	0	0	2.6	6.5
Quin olone-resistant E cali	136	11,776	7.30	0	0	3.3	8.2	19.4
Penicillin-resistant pneumococci	46	1331	18.90	0	5.3	13	24.0	50.0
Cefotaxime/ceftriaxone-resistant pneumococci	33	854	7.50	0	0	3.4	9.6	28.0
Non-ICU Inpatier	nt Areas			Percentile				
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%
MRSA	56	42,502	46.00	25.6	31.9	44.9	52.0	60.8
Methicillin-resistant CNS	53	23,525	65.70	52.2	57.1	65.2	71.1	75.9
Vancomycin-resistant Enterococcus spp	55	32,924	12.00	1.9	3.5	7.1	14.2	18.6
Ciprofloxacin/ofloxacin-resistant Pseudomonas aeruginosa	55	21,302	27.70	13	20.5	27.4	36.8	40.6
Levofloxacin-resistant P aeruginosa	30	10,077	30.50	15.6	21.8	28.7	33.3	44.1
Imipenem-resistant P aeruginosa	53	17,142	12.30	5.6	6.8	10.0	14.4	20.6
Ceftazidime-reisistant P aeruginosa	53	19,587	8.80	1.9	4.0	7.0	11.0	14.1
Piperacillin-resistant P aeruginosa	53	16,828	11.60	3.4	6.5	9.2	14.0	18.3
Cef3-resistant Enterobacter spp	50	7509	21.00	7.7	13.9	20.7	25.7	30.9
Carbapenem-resistant Enterobacter spp	46	5976	1.00	0	0	0	1.2	3.2
Cef3-resistant Klebsiella pneumoniae	55	14,204	5.80	0	0.2	1.5	4.4	14.5
Cef3-resistant Escherichia coli	55	40,751	1.50	0	0	0.6	1.7	3.2
Quin olone-resistant E cali	56	40,694	8.20	0.4	1.8	3.6	7.0	18.9
Penicillin-resistant pneumococci	41	3629	18.20	2.6	5.9	12.0	20.0	31.8
Cefotaxime/ceftriaxone-resistant pneumococci	34	2148	7.60	0	0.9	5.2	10.5	16.3

Table 10. Pooled means and percentiles of the distribution of antimicrobial resistance rates*, by all ICUs combined, non-ICU inpatient units and by outpatients, ICARE/AUR, January 1998 through June 2004

All ICUs com	bined			Percentile					
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%	
MRSA	157	22,899	52.90	20.0	32.7	48.1	60.3	67.9	
Methicillin-resistant CNS	141	13,553	76.60	57.0	69.4	76.3	83.8	88.4	
Vancomycin-resistant Enterococcus spp	140	14.				345347764	24.3	39.2	
Ciprofloxacin/ofloxacin-resistant Pseudomonas aeruginosa	13 Fa	cility loo	rati	on		41.3	51.6		
Levofloxacin-resistant P aeruginosa	68	5	chity lo		OII		40.8	47.7	
Imipenem-resistant P aeruginosa	123	/II ca1	tegory				25.5	38	
Ceftazidime-reisistant P aeruginosa	129	12 Cal	legor y				16.9	23.6	
Piperacillin-resistant P aeruginosa	118	11,640	17.50	2.4	7.5	14.3	19.5	31.4	
Cef3-resistant Enterobacter spp	111	5328	27.70	10.0	17.4	26.1	36.4	47.4	
Carbapenem-resistant Enterobacter spp	93	4663	0.70	0	0	0	0	3.8	
Cef3-resistant Klebsiella pneumoniae	1/9	7529	6.20	0	0	2.0	8.0	20.7	
Cef3-resistant Escherichia coli	140	12,011	1.30	0	0	0	2.6	6.5	
Quin olone-resistant E coli	136	11,776	7.30	0	0	3.3	8.2	19.4	
Penicillin-resistant pneumococci	46	1331	18.90	0	5.3	13	24.0	50.0	
Cefotaxime/ceftriaxone-resistant pneumococci	33	854	7.50	0	0	3.4	9.6	28.0	
Non-ICU Inpatie	nt Areas					Percentile			
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%	
MRSA	56	42,502	46.00	25.6	31.9	44.9	52.0	60.8	
Methicillin-resistant CNS	53	23,525	65.70	52.2	57.1	65.2	71.1	75.9	
Vancomycin-resistant Enterococcus spp	55	32,924	12.00	1.9	3.5	7.1	14.2	18.6	
Ciprofloxacin/ofloxacin-resistant Pseudomonas aeruginosa	55	21,302	27.70	13	20.5	27.4	36.8	40.6	
Levofloxacin-resistant P aeruginosa	30	10,077	30.50	15.6	21.8	28.7	33.3	44.1	
Imipenem-resistant P aeruginosa	53	17,142	12.30	5.6	6.8	10.0	14.4	20.6	
Ceftazidime-reisistant P aeruginosa	53	19,587	8.80	1.9	4.0	7.0	11.0	14.1	
Piperacillin-resistant P aeruginosa	53	16,828	11.60	3.4	6.5	9.2	14.0	18.3	
Cef3-resistant Enterobacter spp	50	7509	21.00	7.7	13.9	20.7	25.7	30.9	
Carbapenem-resistant Enterobacter spp	46	5976	1.00	0	0	0	1.2	3.2	
Cef3-resistant Klebsiella pneumoniae	55	14,204	5.80	0	0.2	1.5	4.4	14.5	
Cef3-resistant Escherichia coli	55	40,751	1.50	0	0	0.6	1.7	3.2	
Quin olone-resistant E cali	56	40,694	8.20	0.4	1.8	3.6	7.0	18.9	
Penicillin-resistant pneumococci	41	3629	18.20	2.6	5.9	12.0	20.0	31.8	
Cefotaxime/ceftriaxone-resistant pneumococci	34	2148	7.60	0	0.9	5.2	10.5	16.3	

Table 10. Pooled means and percentiles of the distribution of antimicrobial resistance rates*, by all ICUs combined, non-ICU inpatient units and by outpatients, ICARE/AUR, January 1998 through June 2004

All ICUs comb	oined				Percentile				
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%	
MRSA	157	22,899	52.90	20.0	32.7	48.1	60.3	67.9	
Methicillin-resistant CNS	141	13,553	76.60	57.0	69.4	76.3	83.8	88.4	
Vancomycin-resistant Enterococcus spp	140	14,140	13.90	0	5	13.6	24.3	39.2	
Ciprofloxacin/ofloxacin-resistant Pseudomonas aeruginosa	134	13,473	34.80	8.3	17.4	29.3	41.3	51.6	
Levofloxacin-resistant P aeruginosa	68	5895	35.30	9.7	18.2	29.1	40.8	47.7	
Imipenem-resistant P aeruginosa	123	11,986	19.10	4.8	8.3	13.2	25.5	38	
Ceftazidime-reisistant P aeruginosa	129	12,805	13.90	0	5	10.8	16.9	23.6	
Piperacillin-resistant P aeruginosa	118	11,640	17.50	2.4	7.5	14.3	19.5	31.4	
Cef3-resistant Enterobocter spp	111	5328	7	100		24.1	244	7.4	
Carbapenem-resistant Enterobacter spp	93	4663	Duo	dn	110	aambina	4:01	3.8	
Cef3-resistant Klebsiella pneumonite	119	7529	Dug	-ur	ug (combina		3.8 0.7	
Cef3-resistant Escherichia coli	140	12,011	1.30	0	0	0	2.6	6.5	
Quin olone-resistant E coli	136	11,776	7.30	0	0	3.3	8.2	19.4	
Penicillin-resistant pneumococci	46	1331	18.90	0	5.3	13	24.0	50.0	
Cefotaxime/ceftriaxone-resistant oneumococci	33	854	7.50	0	0	3.4	9.6	28.0	
Non-ICU Inpatier	nt Areas			Percentile					
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%	
MRSA	56	42,502	46.00	25.6	31.9	44.9	52.0	60.8	
Methicillin-resistant CNS	53	23,525	65.70	52.2	57.1	65.2	71.1	75.9	
Vancomycin-resistant Enterococcus spp	5,8	32,924	12.00	1.9	3.5	7.1	14.2	18.6	
Ciprofloxacin/ofloxacin-resistant Pseudomonas ae uginosa	55	21,302	27.70	13	20.5	27.4	36.8	40.6	
Levofloxacin-resistant P aeruginosa	30	10,077	30.50	15.6	21.8	28.7	33.3	44.1	
Imipenem-resistant P aeruginosa	53	17,142	12.30	5.6	6.8	10.0	14.4	20.6	
Ceftazidime-re isistant P aeruginosa	53	19,587	8.80	1.9	4.0	7.0	11.0	14.1	
Piperacillin-resistant P aeruginosa	53	16,828	11.60	3.4	6.5	9.2	14.0	18.3	
Cef3-resistant Enterobocter spp	50	7509	21.00	7.7	13.9	20.7	25.7	30.9	
Carbapenem-resistant Enterobacter spp	46	5976	1.00	0	0	0	1.2	3.2	
Cef3-resistant Klebsiella pneumoniae	55	14,204	5.80	0	0.2	1.5	4.4	14.5	
Cef3-resistant Escherichia coli	55	40,751	1.50	0	0	0.6	1.7	3.2	
Quin olone-resistant E coli	56	40,694	8.20	0.4	1.8	3.6	7.0	18.9	
Penicillin-resistant pneumococci	41	3629	18.20	2.6	5.9	12.0	20.0	31.8	
Cefotaxime/ceftriaxone-resistant pneumococci	34	2148	7.60	0	0.9	5.2	10.5	16.3	

Table 10. Pooled means and percentiles of the distribution of antimicrobial resistance rates*, by all ICUs combined, non-ICU inpatient units and by outpatients, ICARE/AUR, January 1998 through June 2004

All ICUs comb	oined					Percentile		
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%
MRSA	157	22,899	52.90	20.0	32.7	48.1	60.3	67.9
Methicillin-resistant CNS	141	13,553	76.60	57.0	69.4	76.3	83.8	88.4
Vancomycin-resistant Enterococcus spp	140	14,140	13.90	0	5	13.6	24.3	39.2
Ciprofloxacin/ofloxacin-resistant Pseudomonas aeruginosa	134	13,473	34.80	8.3	17.4	29.3	41.3	51.6
Levofloxacin-resistant Paeruginosa	68	5895	35.30	9.7	18.2	29.1	40.8	47.7
Imipenem-resistant P aeruginosa	123	11,986	19.10	4.8	8.3	13.2	25.5	38
Ceftazidime-reisistant P aeruginosa	129	12,805	13.90	0	5	10.8	16.9	23.6
Piperacillin-resistant P aeruginosa	118	11,640	17.50	2.4	7.5	14.3	19.5	31.4
Cef3-resistant Enterobacter spp	111	5328	27.70	10.0	17.4	26.1	36.4	47.4
Carbapenem-resistant Enterobacter spp	93	4663	0.70	0	0	0	0	3.8
Cef3-resistant Klebsiella pneumoniae	119	7529	6.20	0	0	2.0	8.0	20.7
Cef3-resistant Escherichia coli	140	12,011	1.30	0	0	0	2.6	6.5
Quin olone-resistant E cali	136	11,776	7.30	0	0	3.3	8.2	19.4
Penicillin-resistant pneumococci	46	1331	18.90	0	5.3	13	24.0	50.0
Cefotaxime/ceftriaxone-resistant pneumococci	33	854	7.50	0	0	3.4	9.6	28.0
Non-ICU Inpatier	nt Areas					Percentile		
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%
MRSA	56	42,502	46.00	25.6	31.9	44.9	52.0	60.8
Methicillin-resistant CNS	53	23,525	65.70	52.2	57.1	65.2	711	75.9
Vancomycin-resistant Enterococcus spp	55	32,924	12.00	1.9		Y	4:	.6
Ciprofloxacin/ofloxacin-resistant Pseudomonas aeruginosa	55	21,302	27.70	13	-2	Compara	UVE	.6
Levofloxacin-resistant P aeruginosa	30	10,077	30.50	15.6	2	_		.1
Imipenem-resistant P aeruginosa	53	17,142	12.30	5.6	r	ates		.6
Ceftazidime-reisistant P aeruginosa	53	19,587	8.80	1.9	7.0	7.0	1.1.0	c1
Piperacillin-resistant P aeruginosa	53	16,828	11.60	3.4	6.5	9.2	14.0	18.3
Cef3-resistant Enterobacter spp	50	7509	21.00	7.7	13.9	20.7	25.7	30.9
Carbapenem-resistant Enterobacter spp	46	5976	1.00	0	0	0	1.2	3.2
Cef3-resistant Klebsiella pneumoniae	55	14,204	5.80	0	0.2	1.5	4.4	14.5
Cef3-resistant Escherichia coli	55	40,751	1.50	0	0	0.6	1.7	3.2
Quinolone-resistant E cali	56	40,694	8.20	0.4	1.8	3.6	7.0	18.9
Penicillin-resistant pneumococci	41	3629	18.20	2.6	5.9	12.0	20.0	31.8
Cefotaxime/ceftriaxone-resistant pneumococci	34	2148	7.50	0	0.9	5.2	10.5	16.3

Table 10. Pooled means and percentiles of the distribution of antimicrobial resistance rates*, by all ICUs combined, non-ICU inpatient units and by outpatients, ICARE/AUR, January 1998 through June 2004

All ICUs comb	oined					Percentile		
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%
MRSA	157	22,899	52.90	20.0	32.7	48.1	60.3	67.9
Methicillin-resistant CNS	141	13,553	76.60	57.0	69.4	76.3	83.8	88.4
Vancomycin-resistant Enterococcus spp	140	14,140	13.90	0	5	13.6	24.3	39.2
Ciprofloxacin/ofloxacin-resistant Pseudomonas aeruginosa	134	13,473	34.80	8.3	17.4	29.3	41.3	51.6
Levofloxacin-resistant P aeruginosa	68	5895	35.30	9.7	18.2	29.1	40.8	47.7
Imipenem-resistant P aeruginosa	123	11,986	19.10	4.8	8.3	13.2	25.5	38
Ceftazidime-reisistant P aeruginosa	129	12,805	13.90	0	5	10.8	16.9	23.6
Piperacillin-resistant P aeruginosa	118	11,640	17.50	2.4	7.5	14.3	19.5	31.4
Cef3-resistant Enterobacter spp	111	5328	27.70	10.0	17.4	26.1	36.4	47.4
Carbapenem-resistant Enterobacter spp	93	4663	0.70	0	0	0	0	3.8
Cef3-resistant Klebsiella pneumoniae	119	7529	6.20	0	0	2.0	8.0	20.7
Cef3-resistant Escherichia coli	140	12,011	1.30	0	0	0	2.6	6.5
Quin olone-resistant E cali	136	11,776	7.30	0	0	3.3	8.2	19.4
Penicillin-resistant pneumococci	46	1331	18.90	0	5.3	13	24.0	50.0
Cefotaxime/ceftriaxone-resistant pneumococci	33	854	7.50	0	0	3.4	9.6	28.0
Non-ICU Inpatier	nt Areas			Percentile				
Antimicrobial-resistant pathogen	No. units	No. tested	Pooled mean	10%	25%	50% (median)	75%	90%
MRSA	56	42,502	46.00	25.6	31.9	44.9	52.0	60.8
Methicillin-resistant CNS	53	23,525	65.70	52.2	57.1	65.2	71.1	75.9
Vancomycin-resistant Enterococcus spp	55	32,924	12.00	1.9	3.5	7.1	14.2	18.6
Ciprofloxacin/ofloxacin-resistant Pseudomonas aeruginosa	55	21,302	27.70	13	20.5	27.4	36.8	40.6
Levofloxacin-resistant Paeruginosa	30	10,077	30.50	15.6	21.8	28.7	33.3	44.1
Imipenem-resistant P aeruginosa	53	17,142	12.30	5.6	6.8	10.0	14.4	20.6
Ceftazidime-re isistant P aeruginosa	53	19,587	8.80	1.9	4.0	7.0	11.0	14.1
Piperacillin-resistant P aeruginosa	53	16,828	11.60	3.4	6.5	9.2	14.0	18.3
Cef3-resistant Enterobocter spp	50	7509	21.00	7.7	13.9	20.7	25.7	30.9
Carbapenem-resistant Enterobacter spp	46	5976	1.00	0	0	0	1.2	3.2
Cef3-resistant Klebsiella pneumoniae	55	14,204	5.80	0	0.2	1.5	4.4	14.5
Cef3-resistant Escherichia coli	55	40,751	1.50	0	0	0.6	1.7	3.2
Quin olone-resistant E cali	56	40,694	8.20	0.4	1.8	3.6	7.0	18.9
Penicillin-resistant pneumococci	41	3629	18.20	2.6	5.9	12.0	20.0	31.8
Cefotaxime/ceftriaxone-resistant pneumococci	34	2148	7.60	0	0.9	5.2	10.5	16.3

Table 9. Pooled means and percentiles of the distribution of antimicrobial usage rates (defined daily dose* rates[†]), by non-ICU inpatient areas and various types of ICU, ICARE/AUR, January 1998 through June 2004

Non-ICU Inpatie	ent Areas (n = 74)				Percentile				
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%		
Penicillin group	759,866	60.9	9.2	16.7	28.9	63.8	96.5		
Ampicillin group	1,899,047	152.1	83.2	111.1	141.7	186.3	266.9		
Antips eudomonal penicillins	251,036	20.1	3.1	8.1	16.4	29.0	42.9		
Antistaphylococcal penicillins	245,777	19.7	2.9	5.1	12.5	24.2	35.8		
First-generation cephalosporins	982,573	78.7	43.9	57.4	76.1	106.6	125.1		
Second-generation cephalosporins	368,970	29.6	10.3	16.5	25.3	41.5	54.9		
Third-generation cephalosporins	793,340	63.5	21.9	32.2	53.6	79.5	92.5		
Carbapenem group	85,779	6.9	0.4	1.8	4.7	9.4	17.1		
Aztreonam	34,078	2.7	0.1	0.7	1.8	4.3	6.4		
Fluoroquinolones	1,166,836	93.5	37.9	57.9	91.7	130.3	202.2		
Trimethoprim/sulfamethoxazole	595,248	47.7	5.3	14.8	24.5	39.2	106.3		
Vancomycin (oral)	38,279	3.1	0.1	0.5	1.4	2.5	4.2		
Vancomycin (parenteral)	415,887	33.3	13.1	17.1	24.6	41.0	65.7		
Coronary Car	re Unit (n = 32)		Percentile						
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%		
Penicillin group	4296	30.8	0.0	0.4	7.2	41.7	106.4		
Ampicillin group	12,356	88.5	8.6	44.8	88.3	172.1	245.9		
Antips eudomonal penicillins	4599	32.9	0.0	3.3	20.8	42.8	58.6		
Antistaphylococcal penicillins	3679	26.3	0.0	3.6	12.0	46.2	68.2		
First-generation cephalosporins	6978	50.0	9.0	27.7	36.5	54.4	104.9		
Second-generation cephalosporins	4286	30.7	1.5	7.1	19.8	32.5	42.4		
Third-generation cephalosporins	12,540	89.8	25.1	32.8	73.5	98.0	143.5		
Carbapenem group	1635	11.7	0.0	0.2	6.1	12.1	27.4		
Aztreonam	777	5.6	0.0	0.0	2.0	10.8	14.9		
Fluoroquinolones	12,390	88.7	11.3	27.3	58.9	112.3	214.4		
Trimethoprim/sulfamethoxazole	5585	40.0	0.0	6.0	16.5	43.4	112.8		
Vancomycin (oral)	526	3.8	0.0	0.0	0.0	1.3	7.0		
Vancomycin (parenteral)	7713	55.2	11.2	19.8	36.9	89.3	105.9		

Table 9. Pooled means and percentiles of the distribution of antimicrobial usage rates (defined daily dose* rates[†]), by non-ICU inpatient areas and various types of ICU, ICARE/AUR, January 1998 through June 2004

Non-ICO Inpate	ent Areas (n = 74)		Percentile						
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%		
Penicillin group	759,866	60.9	9.2	16.7	28.9	63.8	96.5		
Ampicillin group	1,899,047	152.1	83.2	111.1	141.7	186.3	266.9		
Antips eudomonal penicillins	251,036	20.1	3.1	8.1	16.4	29.0	42.9		
Antistaphylococcal penicillins	245,777	19.7	2.9	5.1	12.5	24.2	35.8		
First-generation cephalosporins	982,573	78.7	43.9	57.4	76.1	106.6	125.1		
Second-generation cephalosporins	368,970	29.6	10.3	16.5	25.3	41.5	54.9		
Third-generation cephalosporins	793,340	63.5	21.9	32.2	53.6	79.5	92.5		
Carbapenem group	85,779	6.9	0.4	1.8	4.7	9.4	17.1		
Aztreonam	34,078	2.7	0.1	0.7	1.8	4.3	6.4		
Fluoroquinolones	1,166,836	93.5	37.9	57.9	91.7	130.3	202.2		
Trimethoprim/sulfamethoxazole	595,248	47.7	5.3	14.8	24.5	39.2	106.3		
Vancomycin (oral)	38,279	3.1	0.1	0.5	1.4	2.5	4.2		
Vancomycin (parenteral)	415,887	33.3	13.1	17.1	24.6	41.0	65.7		
Coronary Care Unit (n = 32)			1 - 1				~~		
			13.1	1832	Percentile	6562			
		Pooled mean	10%	25%		75%	90%		
Coronary Car	re Unit (n = 32)		2856 Pac	10352	Percentile	7000	7000		
Coronary Car Antimicrobial agent	Ne. DDD	Pooled mean	10%	25%	Percentile 50% (median)	75%	90%		
Coronary Car Antimicrobial agent Penicillin group	No. DDD*	Po oled mean 30.8	10%	25% 0.4	Percentile 50% (median) 7.2	75 %	90% 106.4 245.9		
Antimicrobial agent Penicillin group Ampicillin group	Ne. DDD* 4296 12,356	Pooled mean 30.8 88.5	0.0 8.6	25% 0.4 44.8	Percentile 50% (median) 7.2 88.3	75 % 41.7 172.1	90%		
Antimicrobial agent Penicillin group Ampicillin group Antips eudomonal penicillins Antistaphylococcal penicillins	Ne. DDD+ 4296 12,356 4599	Pooled mean 30.8 88.5 32.9	0.0 8.6 0.0	25% 0.4 44.8 3.3	Percentile 50% (median) 7.2 88.3 20.8	75% 41.7 172.1 42.8	90% 106.4 245.9 58.6 68.2		
Antimicrobial agent Penicillin group Ampicillin group Antips eudomonal penicillins Antistaphylococcal penicillins First-generation cephalosporins	Ne. DDD* 4296 12,356 4599 3679	Pooled mean 30.8 88.5 32.9 26.3	0.0 8.6 0.0 0.0	25% 0.4 44.8 3.3 3.6	Percentile 50% (median) 7.2 88.3 20.8 12.0	75% 41.7 172.1 42.8 46.2	90% 106.4 245.9 58.6 68.2 104.9		
Antimicrobial agent Penicillin group Ampicillin group Antips eudomonal penicillins Antistaphylococcal penicillins First-generation cephalosporins Second-generation cephalosporins	Ne. DDD* 4296 12,356 4599 3679 6978	90 oled mean 30.8 88.5 32.9 26.3 50.0	0.0 8.6 0.0 0.0 9.0	25% 0.4 44.8 3.3 3.6 27.7	Percentile 50% (median) 7.2 88.3 20.8 12.0 36.5	75% 41.7 172.1 42.8 46.2 54.4	90% 106.4 245.9 58.6 68.2 104.9 42.4		
Antimicrobial agent Penicillin group Ampicillin group Antips eudomonal penicillins Antistaphylococcal penicillins First-generation cephalosporins Second-generation cephalosporins Third-generation cephalosporins	4296 12,356 4599 3679 6978 4286	90 oled mean 30.8 88.5 32.9 26.3 50.0 30.7	0.0 8.6 0.0 0.0 9.0 1.5	25% 0.4 44.8 3.3 3.6 27.7 7.1	Percentile 50% (median) 7.2 88.3 20.8 12.0 36.5 19.8	75% 41.7 172.1 42.8 46.2 54.4 32.5	90% 106.4 245.9 58.6 68.2 104.9 42.4 143.5		
Antimicrobial agent Penicillin group Ampicillin group Antips eudomonal penicillins Antistaphylococcal penicillins First-generation cephalosporins Second-generation cephalosporins	4296 12,356 4599 3679 6978 4286 12,540	Pooled mean 30.8 88.5 32.9 26.3 50.0 30.7 89.8	0.0 8.6 0.0 0.0 9.0 1.5 25.1	25% 0.4 44.8 3.3 3.6 27.7 7.1 32.8	Percentile 50% (median) 7.2 88.3 20.8 12.0 36.5 19.8 73.5	75% 41.7 172.1 42.8 46.2 54.4 32.5 98.0	90% 106.4 245.9 58.6 68.2 104.9 42.4 143.5 27.4		
Antimicrobial agent Penicillin group Ampicillin group Antips eudomonal penicillins Antistaphylococcal penicillins First-generation cephalosporins Second-generation cephalosporins Third-generation cephalosporins Carbapenem group Aztreonam	4296 12,356 4599 3679 6978 4286 12,540 1635	90 oled mean 30.8 88.5 32.9 26.3 50.0 30.7 89.8 11.7	0.0 8.6 0.0 0.0 9.0 1.5 25.1	25% 0.4 44.8 3.3 3.6 27.7 7.1 32.8 0.2	Percentile 50% (median) 7.2 88.3 20.8 12.0 36.5 19.8 73.5 6.1	75% 41.7 172.1 42.8 46.2 54.4 32.5 98.0 12.1	90% 106.4 245.9 58.6 68.2 104.9 42.4 143.5 27.4		
Antimicrobial agent Penicillin group Ampicillin group Antips eudomonal penicillins Antistaphylococcal penicillins First-generation cephalosporins Second-generation cephalosporins Third-generation cephalosporins Carbapenem group	4296 12,356 4599 3679 6978 4286 12,540 1635 777	90 oled mean 30.8 88.5 32.9 26.3 50.0 30.7 89.8 11.7 5.6	0.0 8.6 0.0 0.0 9.0 1.5 25.1 0.0	25% 0.4 44.8 3.3 3.6 27.7 7.1 32.8 0.2 0.0	Percentile 50% (median) 7.2 88.3 20.8 12.0 36.5 19.8 73.5 6.1 2.0	75% 41.7 172.1 42.8 46.2 54.4 32.5 98.0 12.1 10.8	90% 106.4 245.9 58.6 68.2 104.9 42.4 143.5 27.4 14.9 214.4		
Antimicrobial agent Penicillin group Ampicillin group Antips eudomonal penicillins Antistaphylococcal penicillins First-generation cephalosporins Second-generation cephalosporins Third-generation cephalosporins Carbapenem group Aztreonam Fluoroquinolones	4296 12,356 4599 3679 6978 4286 12,540 1635 777 12,390	Pooled mean 30.8 88.5 32.9 26.3 50.0 30.7 89.8 11.7 5.6 88.7	0.0 8.6 0.0 0.0 9.0 1.5 25.1 0.0 0.0	25% 0.4 44.8 3.3 3.6 27.7 7.1 32.8 0.2 0.0 27.3	Percentile 50% (median) 7.2 88.3 20.8 12.0 36.5 19.8 73.5 6.1 2.0 58.9	75% 41.7 172.1 42.8 46.2 54.4 32.5 98.0 12.1 10.8 112.3	90% 106.4 245.9 58.6		

Table 9. Pooled means and percentiles of the distribution of antimicrobial usage rates (defined daily dose* rates[†]), by non-ICU inpatient areas and various types of ICU, ICARE/AUR, January 1998 through June 2004

Non-ICU Inpatie	ent Areas (n = 74)		Percentile						
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%		
Penicillin group	759,866	60.9	9.2	16.7	28.9	63.8	96.5		
Ampicillin group	1,899,047	152.1	83.2	111.1	141.7	186.3	266.9		
Antips eudomonal penicillins	251,036	20.1	3.1	8.1	16.4	29.0	42.9		
Antistaphylococcal penicillins	245,777	19.7	2.9	5.1	12.5	24.2	35.8		
First-generation cephalosporins	982,573	78.7	43.9	57.4	76.1	106.6	125.1		
Second-generation cephalosporins	368,970	29.6	10.3	16.5	25.3	41.5	54.9		
Third-generation cephalosporins	793,340	63.5	21.9	32.2	53.6	79.5	92.5		
Carbapenem group	85,779	4 • 4 •	• •1	•	4.7	9.4	17.1		
Aztreonam	34,078 A	Agents with	ı sımıl	lar	1.8	4.3	6.4		
Fluoroquinolones	1.166.836			_	91.7	130.3	202.2		
Trimeth oprim/sulfamethoxazole	595,248 S	pectrum a	re gro	uped	24.5	39.2	106.3		
Vancomycin (oral)	38,279	P	8		1.4	2.5	4.2		
Vancomycin (parenteral)	415,887	33.3	13.1	17.1	24.6	41.0	65.7		
Coronary Car	re Unit (n = 32)		Percentile						
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%		
Penicillin group	4296	30.8	0.0	0.4	7.2	41.7	106.4		
Ampicillin group	2,356	88.5	8.6	44.8	88.3	172.1	245.9		
Antips eudomonal penicillins	4599	32.9	0.0	3.3	20.8	42.8	58.6		
Antistaphylococcal penicillins	3679	26.3	0.0	3.6	12.0	46.2	68.2		
First-generation cephalosporins	6978	50.0	9.0	27.7	36.5	54.4	104.9		
Second-generation cephalosporins	4286	30.7	1.5	7.1	19.8	32.5	42.4		
Third-generation cephalosporins	12,540	89.8	25.1	32.8	73.5	98.0	143.5		
Carbapenem group	1635	11.7	0.0	0.2	6.1	12.1	27.4		
Aztreonam	777	5.6	0.0	0.0	2.0	10.8	14.9		
Fluoroguinolones	12,390	88.7	11.3	27.3	58.9	112.3	214.4		
Trimethoprim/sulfamethoxazole	5585	40.0	0.0	6.0	16.5	43.4	112.8		
Vancomycin (oral)	526	3.8	0.0	0.0	0.0	1.3	7.0		
Vancomycin (parenteral)	7713	55.2	11.2	19.8	36.9	89.3	105.9		

Table 9. Pooled means and percentiles of the distribution of antimicrobial usage rates (defined daily dose* rates[†]), by non-ICU inpatient areas and various types of ICU, ICARE/AUR, January 1998 through June 2004

Non-ICU Inpatient Areas (n = 74)				Percentile				
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%	
Penicillin group	759,866	60.9	9.2	16.7	28.9	63.8	96.5	
Ampicillin group	1,899,047	152.1	83.2	111.1	141.7	186.3	266.9	
Antips eudomonal penicillins	251,036	20.1	3.1	8.1	16.4	29.0	42.9	
Antistaphylococcal penicillins	245,777	19.7	2.9	5.1	12.5	24.2	35.8	
First-generation cephalosporins	982,573	78.7	43.9	57.4	76.1	106.6	125.1	
Second-generation cephalosporins	368,970	29.6	10.3	16.5	25.3	41.5	54.9	
Third-generation cephalosporins	793,340	63.5	21.9	32.2	53.6	79.5	92.5	
Carbapenem group	85,779	6.9	0.4	1.8	4.7	9.4	17.1	
Aztreonam	34,078	2.7	0.1	0.7	1.8	4.3	6.4	
Fluoroquinolones	1,166,836	93.5	37.9	57.9	91.7	130.3	202.2	
Trimeth oprim/sulfamethoxazole	595,248	47.7	300000	1202	1-00/2	12	106.3	
Vancomycin (oral)	38,279	3.1	Numb	or of T	DDD repor	tod 5	4.2	
Vancomycin (parenteral)	415,887	33.3	Mullip		DD Tepor	teu ₀	65.7	
Coronary Ca	re Unit (n = 32)				Percentile			
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%	
Penicillin group	4296	30.8	0.0	0.4	7.2	41.7	106.4	
Ampicillin group	12,356	88.5	8.6	44.8	88.3	172.1	245.9	
Antips eudomonal penicillins	4599	32.9	0.0	3.3	20.8	42.8	58.6	
Antistaphylococcal penicillins	3679	26.3	0.0	3.6	12.0	46.2	68.2	
First-generation cephalosporins	6978	50.0	9.0	27.7	36.5	54.4	104.9	
Second-generation cephalosporins	4286	30.7	1.5	7.1	19.8	32.5	42.4	
Third-generation cephalosporins	12,540	89.8	25.1	32.8	73.5	98.0	143.5	
Carbapenem group	1635	11.7	0.0	0.2	6.1	12.1	27.4	
Aztreonam	777	5.6	0.0	0.0	2.0	10.8	14.9	
Fluoroquinolones	12,390	88.7	11.3	27.3	58.9	112.3	214.4	
[1]	0.0000000000000000000000000000000000000	40.0	0.0	6.0	16.5	43.4	112.8	
Trimethoprim/sulfamethoxazole	5585	70.0	W.W.					
Vancomycin (oral)	526	3.8	0.0	0.0	0.0	1.3	7.0	


Table 9. Pooled means and percentiles of the distribution of antimicrobial usage rates (defined daily dose* rates[†]), by non-ICU inpatient areas and various types of ICU, ICARE/AUR, January 1998 through June 2004

Non-ICU Inpatient Areas (n = 74)			Percentile					
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%	
Penicillin group	759,866	60.9	9.2	16.7	28.9	63.8	96.5	
Ampicillin group	1,899,047	152.1	83.2	111.1	141.7	186.3	266.9	
Antips eudomonal penicillins	251,036	20.1	3.1	8.1	16.4	29.0	42.9	
Antistaphylococcal penicillins	245,777	19.7	2.9	5.1	12.5	24.2	35.8	
First-generation cephalosporins	982,573	78.7	43.9	57.4	76.1	106.6	125.1	
Second-generation cephalosporins	368,970	29.6	10.3	16.5	25.3	41.5	54.9	
Third-generation cephalosporins	793,340	63.5	21.9	32.2	53.6	79.5	92.5	
Carbapenem group	85,779	6.9	0.4	1.8	4.7	9.4	17.1	
Aztreonam	34,078	2.7	0.1	0.7	1.8	4.3	6.4	
Fluoroquinolones	1,166,836	93.5	37.9	57.9	91.7	130.3	202.2	
Trimeth oprim/sulfamethoxazole	595,248	47.7	5.3	14.8	24.5	39.2	106.3	
Vancomycin (oral)	38,279	3.1	0.1	0.5	1.4	2.5	4.2	
Vancomycin (parenteral)	415,887	33.3	13.1	17.1	24.6	41.0	65.7	
Coronary Care Unit (n = 32)					Percentile			
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%	
Penicillin group	4296	30.8				_	106.4	
Ampicillin group	12,356	88.5	Pooled mean for each					
Antips eudomonal penicillins	4599	32.9						
Antistaphylococcal penicillins	3679	26.3	∣anti	micro	bial agent		68.2	
First-generation cephalosporins	6978	50.0	302202				104.9	
Second-generation cephalosporins	4286	30.7	grou	ın			42.4	
Third-generation cephalosporins	12,540	89.8	810	"P			143.5	
Carbapenem group	1635	11.7	0.0	0.2	6.1	12.1	27.4	
Aztreonam	777	5.6	0.0	0.0	2.0	10.8	14.9	
Fluoroquinolones	12,390	88.7	11.3	27.3	58.9	112.3	214.4	
Trimethoprim/sulfamethoxazole	5585	40.0	0.0	6.0	16.5	43.4	112.8	
Vancomycin (oral)	526	3.8	0.0	0.0	0.0	1.3	7.0	

Table 9. Pooled means and percentiles of the distribution of antimicrobial usage rates (defined daily dose* rates[†]), by non-ICU inpatient areas and various types of ICU, ICARE/AUR, January 1998 through June 2004

Non-ICU Inpatient Areas (n = 74)			Percentile					
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%	
Penicillin group	759,866	60.9	9.2	16.7	28.9	63.8	96.5	
Ampicillin group	1,899,047	152.1	83.2	111.1	141.7	186.3	266.9	
Antips eudomonal penicillins	251,036	20.1	3.1	8.1	16.4	29.0	42.9	
Antistaphylococcal penicillins	245,777	19.7	2.9	5.1	12.5	24.2	35.8	
First-generation cephalosporins	982,573	78.7	43.9	57.4	76.1	106.6	125.1	
Second-generation cephalosporins	368,970	29.6	10.3	16.5	25.3	41.5	54.9	
Third-generation cephalosporins	793,340	63.5	21.9	32.2	53.6	79.5	92.5	
Carbapenem group	85,779	6.9	0.4	1.8	4.7	9.4	17.1	
Aztreonam	34,078	2.7	0.1	0.7	1.8	4.3	6.4	
Fluoroquinolones	1,166,836	93.5	37.9	57.9	91.7	130.3	202.2	
Trimeth op rim/s ulfamethoxazole	595,248	47.7	5.3	14.8	24.5	39.2	106.3	
Vancomycin (oral)	38,279	3.1	0.1	0.5	1.4	2.5	4.2	
Vancomycin (parenteral)	415,887	33.3	13.1	17.1	24.6	41.0	65.7	
Coronary Care Unit (n = 32)			Percentile					
Antimicrobial agent	No. DDD*	Pooled mean	10%	25%	50% (median)	75%	90%	
Penicillin group	4296	30.8	0.0	0.4	7.2	41.7	106.4	
Ampicillin group	12,356	88.5	8.6	44.8	88.3	172.1	245.9	
Antips eudomonal penicillins	4599	32.9	0.0	3.3	20.8	42.8	58.6	
Antistaphylococcal penicillins	3679	26.3	0.0	3.6	12.0	46.2	68.2	
First-generation cephalosporins	6978	50.0	9.0	27.7	36.5	54.4	104.9	
Second-generation cephalosporins	4286	30.7	1.5	7.1	19.8	32.5	42.4	
Third-generation cephalosporins	12,540	89.8	25.1	32.8	73.5	98.0	143.5	
Carbapenem group	1635	11.7	0.0	0.2	6.1	12.1	27.4	
Aztreonam	777	5.6	0.0	0.0	2.0	10.8	14.9	
Fluoroguinolones	12,390	88.7	11.3	27.3	58.9	112.3	214.4	
Trimethoprim/sulfamethoxazole	5585	40.0	0.0	6.0	16.5	43.4	112.8	
Vancomycin (oral)	526	3.8	0.0	0.0	0.0	1.3	7.0	
Vancomycin (parenteral)	7713	55.2	11.2	19.8	36.9	89.3	105.9	

Using the Data



- Detect emergence of new resistant organisms or changes in resistance patterns in your facility
- Identify changes that may correlate with
 - Staffing changes
 - Staff education
 - Antibiotic prescribing practices
- Compare your rates with NHSN rates

Using the Data

- Helpful in getting administrative support for infection control intervention activities
- Correlate data with pathogens identified in infections (e.g., VRE increasing rates and increasing incidence of CLABSI with VRE pathogen)

Using the Data

 Provide clinicians with information to guide antimicrobial prescribing practices







Questions?

http://www.cdc.gov/ncidod/dhqp/nhsn_members.html

SAFER · HEALTHIER · PEOPLE™